

Math 537 - Ordinary Differential Equations

Lecture Notes – Method of Frobenius

Joseph M. Mahaffy,
(jmahaffy@sdsu.edu)

Department of Mathematics and Statistics
Dynamical Systems Group
Computational Sciences Research Center
San Diego State University
San Diego, CA 92182-7720

<http://jmahaffy.sdsu.edu>

Fall 2019

Outline

- 1 **Definitions**
 - Legendre's Equation

- 2 **Cauchy-Euler Equation**
 - Distinct Roots
 - Equal Roots
 - Complex Roots

- 3 **Method of Frobenius**
 - Distinct Roots, $r_1 - r_2 \neq N$
 - Repeated Roots, $r_1 = r_2 = r$
 - Roots Differing by an Integer, $r_1 - r_2 = N$

Definitions

Consider the 2^{nd} order linear differential equation:

$$P(x)y'' + Q(x)y' + R(x)y = F(x). \quad (1)$$

Definition (Ordinary and Singular Points)

x_0 is an **ordinary point** of Eqn. (1) if $P(x_0) \neq 0$ and $Q(x)/P(x)$, $R(x)/P(x)$, and $F(x)/P(x)$ are **analytic** at x_0 .

x_0 is a **singular point** of Eqn. (1) if x_0 is not an **ordinary point**.

The previous **ODEs** solved by **power series** methods have centered around $x_0 = 0$, when this is an **ordinary point**.

In an interval about a **singular point**, the solutions of Eqn. (1) can exhibit behavior different from **power series solutions** for Eqn. (1) near an **ordinary point**.

If $x_0 = 0$, then these solutions may behave like $\ln(x)$ or x^{-n} near x_0 .

Definitions

We concentrate on the *homogeneous 2nd order linear differential equation* to better understand behavior of the solution near a *singular point*:

$$P(x)y'' + Q(x)y' + R(x)y = 0. \quad (2)$$

Definition (Regular and Irregular Singular Points)

If x_0 is a *singular point* of Eqn. (2), then x_0 is a *regular singular point* provided the functions:

$$(x - x_0) \frac{Q(x)}{P(x)} \quad \text{and} \quad (x - x_0)^2 \frac{R(x)}{P(x)}$$

are *analytic* at x_0 .

A *singular point* that is **not regular** is said to be an *irregular singular point*.

If Eqn. (2) has a *regular singular point* at x_0 , then it is **possible** that **no power series** solution exists of the form:

$$y(x) = \sum_{n=0}^{\infty} a_n (x - x_0)^n.$$

Bessel's Equation

Example: Consider *Bessel's equation of order ν* :

$$x^2 y'' + xy' + (x^2 - \nu^2)y = 0,$$

where $P(x) = x^2$, $Q(x) = x$, and $R(x) = x^2 - \nu^2$.

It is clear that $x = 0$ is a *singular point*.

We see that

$$\lim_{x \rightarrow 0} x \frac{Q(x)}{P(x)} = 1 \quad \text{and} \quad \lim_{x \rightarrow 0} x^2 \frac{R(x)}{P(x)} = \lim_{x \rightarrow 0} (x^2 - \nu^2) = -\nu^2,$$

which are both finite, so *analytic*.

It follows that $x_0 = 0$ is a *regular singular point*.

Any other value of x_0 for *Bessel's equation* gives an *ordinary point*.

Legendre's Equation

Example: Consider *Legendre's equation*:

$$(1 - x^2)y'' - 2xy' + \alpha(\alpha + 1)y = 0,$$

where $P(x) = (1 - x^2)$, $Q(x) = -2x$, and $R(x) = \alpha(\alpha + 1)$.

It is clear that $x = \pm 1$ are *singular points*.

We see that

$$\lim_{x \rightarrow 1} (x - 1) \frac{Q(x)}{P(x)} = \lim_{x \rightarrow 1} (x - 1) \frac{-2x}{(1 - x^2)} = \lim_{x \rightarrow 1} \frac{2x}{1 + x} = 1, \quad \text{and}$$

$$\lim_{x \rightarrow 1} (x - 1)^2 \frac{R(x)}{P(x)} = \lim_{x \rightarrow 1} (x - 1)^2 \frac{\alpha(\alpha + 1)}{(1 - x^2)} = \lim_{x \rightarrow 1} (x - 1) \frac{-\alpha(\alpha + 1)}{1 + x} = 0,$$

which are both finite, so *analytic*.

It follows that $x_0 = 1$ is a *regular singular point*, and a similar argument shows that $x_0 = -1$ is a *regular singular point*.

Legendre's Equation

Any other value of x_0 for *Legendre's equation* gives an *ordinary point*, so $x_0 = 0$ is an *ordinary point*, and we seek power series solutions:

$$y(x) = \sum_{n=0}^{\infty} a_n x^n, \quad y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}, \quad \text{and} \quad y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

These are inserted into the *Legendre Equation* to give:

$$(1-x^2) \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} - 2x \sum_{n=1}^{\infty} n a_n x^{n-1} + \alpha(\alpha+1) \sum_{n=0}^{\infty} a_n x^n = 0$$

The first two sums could start their index at $n = 0$ without changing anything, so this expression is easily changed by multiplying by x or x^2 and shifting the index to:

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} n(n-1) a_n x^n - 2 \sum_{n=0}^{\infty} n a_n x^n + \alpha(\alpha+1) \sum_{n=0}^{\infty} a_n x^n = 0.$$

Legendre's Equation

Collecting coefficients gives:

$$\sum_{n=0}^{\infty} \left[(n+2)(n+1)a_{n+2} - (n(n-1) + 2n - \alpha(\alpha+1))a_n \right] x^n = 0$$

or

$$\sum_{n=0}^{\infty} \left[(n+2)(n+1)a_{n+2} - (n(n+1) - \alpha(\alpha+1))a_n \right] x^n = 0.$$

The previous expression gives the **recurrence relation**:

$$a_{n+2} = \frac{n(n+1) - \alpha(\alpha+1)}{(n+2)(n+1)} a_n \quad \text{for } n = 0, 1, ..$$

Properties of power series give a_0 and a_1 as arbitrary with $y(0) = a_0$ and $y'(0) = a_1$.

Legendre's Equation

The **recurrence relation** shows that all the even coefficients, a_{2n} , depend only on a_0 , while all odd coefficients, a_{2n+1} , depend only on a_1 , so all solutions have the form:

$$y(x) = a_0 y_1(x) + a_1 y_2(x),$$

where $y_1(x)$ has only even powers of x and $y_2(x)$ has only odd powers of x .

From the **recurrence relation** it is clear that any integer value of $\alpha = 0, 1, 2, \dots$ results in coefficients $a_{\alpha+2} = a_{\alpha+4} = \dots = a_{\alpha+2k} = 0$ for $k = 1, 2, \dots$

This results in one solution being an **α -degree polynomial**, which is valid for all x .

The other solution remains an **infinite series**.

If α is not an integer, then both **linearly independent solutions** are **infinite series**.

The **polynomial solution** converges for all x , while the **infinite series solution** converges for $|x| < 1$ using the **ratio test**.

Cauchy-Euler Equation

Cauchy-Euler Equation (Also, **Euler Equation**): Consider the differential equation:

$$L[y] = t^2 y'' + \alpha t y' + \beta y = 0,$$

where α and β are constants.

Assume $t > 0$ and attempt a solution of the form

$$y(t) = t^r.$$

Note that t^r may not be defined for $t < 0$.

The result is

$$\begin{aligned} L[t^r] &= t^2(r(r-1)t^{r-2}) + \alpha t(rt^{r-1}) + \beta t^r \\ &= t^r[r(r-1) + \alpha r + \beta] = 0. \end{aligned}$$

Thus, obtain quadratic equation

$$F(r) = r(r-1) + \alpha r + \beta = 0.$$

Cauchy-Euler Equation

Cauchy-Euler Equation: The quadratic equation

$$F(r) = r(r - 1) + \alpha r + \beta = 0$$

has roots

$$r_1, r_2 = \frac{-(\alpha - 1) \pm \sqrt{(\alpha - 1)^2 - 4\beta}}{2}.$$

This is very similar to our **constant coefficient homogeneous DE**.

Real, Distinct Roots: If $F(r) = 0$ has real roots, r_1 and r_2 , with $r_1 \neq r_2$, then the **general solution** of

$$L[y] = t^2 y'' + \alpha y' + \beta y = 0,$$

is

$$y(t) = c_1 t^{r_1} + c_2 t^{r_2}, \quad t > 0.$$

Cauchy-Euler Equation

Example: Consider the equation

$$2t^2y'' + 3ty' - y = 0.$$

By substituting $y(t) = t^r$, we have

$$t^r[2r(r-1) + 3r - 1] = t^r(2r^2 + r - 1) = t^r(2r-1)(r+1) = 0.$$

This has the real roots $r_1 = -1$ and $r_2 = \frac{1}{2}$, giving the **general solution**

$$y(t) = c_1t^{-1} + c_2\sqrt{t}, \quad t > 0.$$

Cauchy-Euler Equation

Equal Roots: If $F(r) = (r - r_1)^2 = 0$ has r_1 as a double root, there is one solution, $y_1(t) = t^{r_1}$.

Need a second linearly independent solution.

Note that not only $F(r_1) = 0$, but $F'(r_1) = 0$, so consider

$$\begin{aligned}\frac{\partial}{\partial r} L[t^r] &= \frac{\partial}{\partial r} [t^r F(r)] = \frac{\partial}{\partial r} [t^r (r - r_1)^2] \\ &= (r - r_1)^2 t^r \ln(t) + 2(r - r_1) t^r.\end{aligned}$$

Also,

$$\frac{\partial}{\partial r} L[t^r] = L \left[\frac{\partial}{\partial r} (t^r) \right] = L[t^r \ln(t)].$$

Evaluating these at $r = r_1$ gives

$$L[t^{r_1} \ln(t)] = 0.$$

Cauchy-Euler Equation

Equal Roots: For $F(r) = (r - r_1)^2 = 0$, where r_1 is a double root, then the differential equation

$$L[y] = t^2 y'' + \alpha y' + \beta y = 0,$$

was shown to satisfy

$$L[t^{r_1}] = 0 \quad \text{and} \quad L[t^{r_1} \ln(t)] = 0.$$

It follows that the **general solution** is

$$y(t) = (c_1 + c_2 \ln(t))t^{r_1}.$$

Cauchy-Euler Equation

Example: Consider the equation

$$t^2 y'' + 5ty' + 4y = 0.$$

By substituting $y(t) = t^r$, we have

$$t^r [r(r-1) + 5r + 4] = t^r (r^2 + 4r + 4) = t^r (r+2)^2 = 0.$$

This only has the real root $r_1 = -2$, which gives **general solution**

$$y(t) = (c_1 + c_2 \ln(t))t^{-2}, \quad t > 0.$$

Cauchy-Euler Equation

Complex Roots: Assume $F(r) = 0$ has $r = \mu \pm i\nu$ as complex roots, the solutions are still $y(t) = t^r$.

However,

$$t^r = e^{(\mu+i\nu)\ln(t)} = t^\mu[\cos(\nu \ln(t)) + i \sin(\nu \ln(t))].$$

As before, we obtain the two linearly independent solutions by taking the real and imaginary parts, so the **general solution** is

$$y(t) = t^\mu[c_1 \cos(\nu \ln(t)) + c_2 \sin(\nu \ln(t))].$$

Cauchy-Euler Equation

Example: Consider the equation

$$t^2 y'' + t y' + y = 0.$$

By substituting $y(t) = t^r$, we have

$$t^r [r(r-1) + r + 1] = t^r (r^2 + 1) = 0.$$

This has the complex roots $r = \pm i$ ($\mu = 0$ and $\nu = 1$), which gives the **general solution**

$$y(t) = c_1 \cos(\ln(t)) + c_2 \sin(\ln(t)), \quad t > 0.$$

Regular Singular Problem

1

Regular Singular Point: Consider the equation

$$P(x)y'' + Q(x)y' + R(x)y = 0,$$

and without loss of generality assume that it has a **regular singular point** at $x_0 = 0$.

This implies that $xQ(x)/P(x) = p(x)$ and $x^2R(x)/P(x) = q(x)$ are **analytic** at $x = 0$, so

$$p(x) = \sum_{n=0}^{\infty} p_n x^n \quad \text{and} \quad q(x) = \sum_{n=0}^{\infty} q_n x^n,$$

are convergent series for some interval $|x| < \rho$ with $\rho > 0$.

This gives the equation:

$$L[y] = x^2y'' + xp(x)y' + q(x)y = 0.$$

Regular Singular Problem

Regular Singular Problem: Since the $p(x)$ and $q(x)$ are *analytic* at $x = 0$, the *second order linear equation* can be written:

$$L[y] = x^2 y'' + x \left(\sum_{n=0}^{\infty} p_n x^n \right) y' + \left(\sum_{n=0}^{\infty} q_n x^n \right) y = 0. \quad (3)$$

Note that if $p_n = q_n = 0$ for $n = 1, 2, \dots$ with

$$p_0 = \lim_{x \rightarrow 0} \frac{xQ(x)}{P(x)} \quad \text{and} \quad q_0 = \lim_{x \rightarrow 0} \frac{x^2 R(x)}{P(x)},$$

then the *second order linear equation* becomes the *Cauchy-Euler equation*:

$$x^2 y'' + x p_0 y' + q_0 y = 0.$$

Method of Frobenius

1

Method of Frobenius: Since the **regular singular problem** starts with the *zeroth order terms* in the coefficients being similar to the *Cauchy-Euler equation*, this suggests looking for solutions with terms of x^r .

As with the *Cauchy-Euler equation*, we consider $x > 0$ with the case $x < 0$ handled by a change of variables $x = -\xi$ with $\xi > 0$.

The **Method of Frobenius** seeks solutions of the form:

$$y(x) = x^r (a_0 + a_1 x + \cdots + a_n x^n + \cdots) = x^r \sum_{n=0}^{\infty} a_n x^n = \sum_{n=0}^{\infty} a_n x^{r+n}.$$

- 1 What values of r give a solution to (3) in the above form?
- 2 What is the *recurrence relation* for the a_n ?
- 3 What is the radius of convergence for the above series?

Method of Frobenius

2

For the **regular singular problem**

$$x^2 y'' + x \left(\sum_{n=0}^{\infty} p_n x^n \right) y' + \left(\sum_{n=0}^{\infty} q_n x^n \right) y = 0,$$

we seek a solutions of the form: $y(x) = \sum_{n=0}^{\infty} a_n x^{r+n}$, so

$$y' = \sum_{n=0}^{\infty} a_n (r+n) x^{r+n-1} \quad \text{and} \quad y'' = \sum_{n=0}^{\infty} a_n (r+n)(r+n-1) x^{r+n-2}.$$

Thus,

$$\begin{aligned} \sum_{n=0}^{\infty} a_n (r+n)(r+n-1) x^{r+n} &+ \left(\sum_{n=0}^{\infty} p_n x^n \right) \sum_{n=0}^{\infty} a_n (r+n) x^{r+n} \\ &+ \left(\sum_{n=0}^{\infty} q_n x^n \right) \sum_{n=0}^{\infty} a_n x^{r+n} = 0. \end{aligned}$$

Method of Frobenius

Indicial Equation: In the previous equation, we examine the lowest power of x , so $n = 0$.

This gives

$$a_0 x^r (r(r-1) + p_0 r + q_0) = 0.$$

For $a_0 \neq 0$, we obtain the *indicial equation*, which came from solving the *Cauchy-Euler equation*:

$$F(r) = r(r-1) + p_0 r + q_0 = 0,$$

which is a *quadratic equation*.

The form of the solution of the *Cauchy-Euler equation* depended on the values of r for the *indicial equation*, which in turn affects the factor x^r multiplying our *power series solution*.

Distinct Roots, $r_1 - r_2 \neq N$

1

The *Method of Frobenius* breaks into **3** cases, depending on the roots of the *indicial equation*.

Case 1. Distinct roots not differing by an integer, $r_1 - r_2 \neq N$.

For this case, a basis for the solution of the *regular singular problem* satisfies:

$$y_1(x) = x^{r_1}(a_0 + a_1x + \cdots + a_nx^n + \dots) = x^{r_1} \sum_{n=0}^{\infty} a_n x^n$$

and

$$y_2(x) = x^{r_2}(b_0 + b_1x + \cdots + b_nx^n + \dots) = x^{r_2} \sum_{n=0}^{\infty} b_n x^n$$

with these solutions converging for at least $|x| < \rho$, where ρ is the radius of convergence for $p(x)$ and $q(x)$.

Distinct Roots, $r_1 - r_2 \neq N$

2

Example: Consider the *regular singular problem* given by

$$4xy'' + 2y' + y = 0,$$

where $x = 0$ is a *regular singular point*.

From our definitions before we have $p(x) = \frac{1}{2}$ and $q(x) = \frac{x}{4}$, which implies that $p_0 = \frac{1}{2}$ and $q_0 = 0$.

Since p and q have convergent power series for all x , the solutions will converge for $|x| < \infty$.

The *indicial equation* is given by:

$$r(r-1) + \frac{1}{2}r = r\left(r - \frac{1}{2}\right) = 0,$$

so $r_1 = 0$ and $r_2 = \frac{1}{2}$.

Distinct Roots, $r_1 - r_2 \neq N$

Example: We multiply our example by x and continue with

$$4x^2y'' + 2xy' + xy = 0,$$

trying a solution of the form: $y = \sum_{n=0}^{\infty} a_n x^{n+r}$.

Differentiating y and entering into the equation gives:

$$4 \sum_{n=0}^{\infty} a_n (r+n)(r+n-1)x^{r+n} + 2 \sum_{n=0}^{\infty} a_n (r+n)x^{r+n} + \sum_{n=1}^{\infty} a_{n-1} x^{r+n} = 0,$$

shifting the last index to match powers of x .

Note that when $n = 0$, we have

$$a_0 [4r(r-1) + 2r] = 2a_0 r(2r-1) = 0,$$

which is an alternate way to obtain the *indicial equation*.

Distinct Roots, $r_1 - r_2 \neq N$

Example: For $n \geq 1$, we match powers of x , so

$$[4(r+n)(r+n-1) + 2(r+n)]a_n + a_{n-1} = 0.$$

From this we obtain the *recurrence relation*:

$$a_{n+1} = \frac{-a_n}{(2n+2r+2)(2n+2r+1)}, \quad \text{for } n = 0, 1, \dots$$

First Solution: Let $r = r_1 = 0$, then the *recurrence relation* becomes:

$$a_{n+1} = \frac{-a_n}{(2n+2)(2n+1)}, \quad \text{for } n = 0, 1, \dots,$$

so

$$a_1 = -\frac{a_0}{2 \cdot 1}, \quad a_2 = -\frac{a_1}{4 \cdot 3} = \frac{a_0}{4!}, \quad a_3 = -\frac{a_2}{6 \cdot 5} = -\frac{a_0}{6!}.$$

Thus,

$$a_n = \frac{(-1)^n}{(2n)!} a_0.$$

Distinct Roots, $r_1 - r_2 \neq N$

Example: Thus, the first solution is:

$$y_1(x) = a_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^n.$$

Second Solution: Let $r = r_2 = \frac{1}{2}$, then the *recurrence relation* becomes:

$$b_{n+1} = \frac{-b_n}{(2n+3)(2n+2)}, \quad \text{for } n = 0, 1, \dots,$$

so

$$b_1 = -\frac{b_0}{3 \cdot 2}, \quad b_2 = -\frac{b_1}{5 \cdot 4} = \frac{b_0}{5!}, \quad b_3 = -\frac{b_2}{7 \cdot 6} = -\frac{b_0}{7!}.$$

Thus,

$$b_n = \frac{(-1)^n}{(2n+1)!} b_0.$$

Thus, the second solution is:

$$y_2(x) = b_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{n+\frac{1}{2}}.$$

Distinct Roots, $r_1 - r_2 \neq N$

Example: The series we see as solutions to this problem have similarities with series for **cosine** and **sine**.

Specifically, a change of variables gives:

$$y_1(x) = a_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^n = a_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} (x^{\frac{1}{2}})^{2n} = a_0 \cos(\sqrt{x}).$$

Similarly, the second linearly independent solution satisfies:

$$y_2(x) = b_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{n+\frac{1}{2}} = b_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} (x^{\frac{1}{2}})^{2n+1} = b_0 \sin(\sqrt{x}).$$

Thus, we could write the general solution as

$$y(x) = a_0 \cos(\sqrt{x}) + b_0 \sin(\sqrt{x}),$$

which can readily be shown satisfies the **ODE** in this example:

$$4x^2 y'' + 2xy' + xy = 0.$$

Repeated Roots, $r_1 = r_2 = r$

1

Case 2. Repeated roots, $r_1 = r_2 = r$.

For this case, a basis for the solution of the *regular singular problem* satisfies:

$$y_1(x) = x^r (a_0 + a_1x + \cdots + a_nx^n + \cdots) = x^r \sum_{n=0}^{\infty} a_n x^n$$

and

$$y_2(x) = y_1(x) \ln(x) + x^r (b_1x + \cdots + b_nx^n + \cdots) = y_1(x) \ln(x) + x^r \sum_{n=1}^{\infty} b_n x^n$$

with these solutions converging for at least $|x| < \rho$, where ρ is the radius of convergence for $p(x)$ and $q(x)$.

Repeated Roots, $r_1 = r_2 = r$

The form of the second solution is found in a manner similar to solving the *Cauchy-Euler equation*.

The first solution is found as before with:

$$y_1(x) = \phi(r, x) = x^r \sum_{n=0}^{\infty} a_n(r)x^n,$$

where the coefficients $a_n(r)$ are determined by a *recurrence relation* with the values of r found from the *indicial equation*

$$F(r) = r(r - 1) + p_0r + q_0 = 0.$$

Our *regular singular problem* was $L[y] = x^2y'' + xp(x)y' + q(x)y = 0$, which with our first solution and the power series for $p(x)$ and $q(x)$ gives

$$L[\phi](r, x) = x^r a_0 F(r) + \sum_{n=1}^{\infty} \left[a_n F(r + n) + \sum_{k=0}^{n-1} a_k [(r + k)p_{n-k} + q_{n-k}] \right] x^{r+n} = 0,$$

where the second sum comes from multiplying the infinite series and collecting terms.

Repeated Roots, $r_1 = r_2 = r$

Assuming $F(r+n) \neq 0$, the *recurrence relation* for the coefficients as a function of r satisfies:

$$a_n(r) = -\frac{\sum_{k=0}^{n-1} a_k[(r+k)p_{n-k} + q_{n-k}]}{F(r+n)}, \quad n \geq 1.$$

Selecting these coefficients reduces our power series solution to:

$$L[\phi](r, x) = x^r a_0 F(r),$$

where $F(r) = (r - r_1)^2$ for our repeated root, so $L[\phi](r_1, x) = 0$, since our first solution is:

$$y_1(x) = \phi(r_1, x) = x^{r_1} \sum_{n=0}^{\infty} a_n(r_1) x^n,$$

Significantly, we have

$$L\left[\frac{\partial \phi}{\partial r}\right](r_1, x) = a_0 \left. \frac{\partial}{\partial r} [x^r (r - r_1)^2] \right|_{r=r_1} = a_0 [(r - r_1)^2 x^r \ln(x) + 2(r - r_1)x^r] \Big|_{r=r_1} = 0,$$

so $\frac{\partial \phi}{\partial r}(r_1, x)$ is also a solution to our problem.

Repeated Roots, $r_1 = r_2 = r$

Since $\frac{\partial \phi}{\partial r}(r_1, x)$ is a solution and our first solution is:

$$y_1(x) = \phi(r_1, x) = x^{r_1} \sum_{n=0}^{\infty} a_n(r_1)x^n,$$

we obtain the second solution:

$$\begin{aligned} y_2(x) &= \left. \frac{\partial \phi(r, x)}{\partial r} \right|_{r=r_1} = \left. \frac{\partial}{\partial r} \left[x^r \sum_{n=0}^{\infty} a_n(r)x^n \right] \right|_{r=r_1} \\ &= (x^{r_1} \ln(x)) \sum_{n=0}^{\infty} a_n(r_1)x^n + x^{r_1} \sum_{n=1}^{\infty} a'_n(r_1)x^n \\ &= y_1(x) \ln(x) + x^{r_1} \sum_{n=1}^{\infty} a'_n(r_1)x^n, \quad x > 0, \end{aligned}$$

where

$$a_n(r) = -\frac{\sum_{k=0}^{n-1} a_k[(r+k)p_{n-k} + q_{n-k}]}{F(r+n)}, \quad n \geq 1.$$

Bessel's Equation Order Zero

1

Bessel's Equation Order Zero satisfies:

$$x^2 y'' + xy' + x^2 y = 0,$$

where $x = 0$ is a *regular singular point*.

From our definitions before we have $p(x) = 1$ and $q(x) = x^2$, which implies that $p_0 = 1$ and $q_0 = 0$.

Since p and q have convergent power series for all x , the solutions will converge for $|x| < \infty$.

The *indicial equation* is given by:

$$r(r - 1) + r = r^2 = 0,$$

so $r_1 = r_2 = 0$, *repeated root*.

Bessel's Equation Order Zero

With *Bessel's equation order zero*,

$$x^2 y'' + xy' + x^2 y = 0,$$

we try a solution of the form: $y = \sum_{n=0}^{\infty} a_n x^{n+r}$.

Differentiating y and entering into the equation gives:

$$\sum_{n=0}^{\infty} a_n (n+r)(n+r-1)x^{n+r} + \sum_{n=0}^{\infty} a_n (n+r)x^{n+r} + \sum_{n=2}^{\infty} a_{n-2} x^{n+r} = 0,$$

shifting the last index to match powers of x .

From the same powers of x , $(n+r)^2 a_n + a_{n-2} = 0$, which for $r = 0$ gives the *recurrence relation*:

$$a_n = \frac{-a_{n-2}}{n^2}, \quad \text{for } n = 2, 3, 4, \dots,$$

with a_0 arbitrary and $a_1 = 0$.

Bessel's Equation Order Zero

The *recurrence relation* shows that the **odd powers** of x all vanish.

Letting $n = 2m$ in the *recurrence relation* gives:

$$a_{2m} = \frac{-a_{2m-2}}{(2m)^2}, \quad \text{for } n = 1, 2, 3, \dots,$$

so

$$a_2 = -\frac{a_0}{2^2}, \quad a_4 = -\frac{a_2}{(2 \cdot 2)^2} = \frac{a_0}{2^4 \cdot 2^2}, \quad a_6 = -\frac{a_4}{(2 \cdot 3)^2} = -\frac{a_0}{2^6(3 \cdot 2)^2}, \quad \dots$$

In general, we have


$$a_{2m} = \frac{(-1)^m a_0}{2^{2m}(m!)^2}, \quad m = 1, 2, 3, \dots$$

The *first solution* becomes:

$$y_1(x) = a_0 J_0(x) = a_0 \sum_{m=0}^{\infty} \frac{(-1)^m x^{2m}}{2^{2m}(m!)^2}.$$

Bessel's Equation Order Zero

This *first solution* gives the *Bessel function of the first kind of order zero*, $J_0(x)$. Below shows some polynomial approximations from the partial sums of the series solution.



Bessel's Equation Order Zero

Since the *Bessel's equation order zero* has only the **repeated root** $r = 0$ from the *indicial equation*, the second solution has the form:

$$y_2(x) = y_1(x) \ln(x) + x^r \sum_{n=1}^{\infty} b_n x^n = \ln(x) J_0(x) + \sum_{n=1}^{\infty} b_n x^n.$$

One technique to solve for this second solution is to substitute into *Bessel's equation* and solve for the coefficients, b_n .

Alternately, we use our results in deriving this form of the second solution, where we found that the coefficients satisfied:

$$b_n = a'_n(r), \quad \text{where} \quad a_n(r) = -\frac{a_{n-2}(r)}{(n+r)^2},$$

evaluated at $r = 0$ based on the *recurrence relation* for coefficients of the first solution.

Bessel's Equation Order Zero

From the formula deriving the *recurrence relation*, we find $(r + 1)^2 a_1(r) = 0$, so not only $a_1(0) = 0$, but $a_1'(0) = 0$.

It follows from the *recurrence relation* that

$$a_3'(0) = a_5'(0) = \cdots = a_{2k+1}'(0) = \cdots = 0.$$

The *recurrence relation* gives:

$$a_{2m}(r) = -\frac{a_{2m-2}(r)}{(2m+r)^2}, \quad m = 1, 2, 3, \dots$$

Hence,

$$a_2(r) = -\frac{a_0}{(2+r)^2},$$

$$a_4(r) = -\frac{a_2(r)}{(4+r)^2} = \frac{a_0}{(4+r)^2(2+r)^2},$$

$$a_6(r) = -\frac{a_4(r)}{(6+r)^2} = \frac{a_0}{(6+r)^2(4+r)^2(2+r)^2},$$

$$a_{2m}(r) = \frac{(-1)^m a_0}{(2m+r)^2(2m-2+r)^2 \cdots (4+r)^2(2+r)^2}, \quad m = 1, 2, 3, \dots$$

Bessel's Equation Order Zero

Note that if

$$f(x) = (x - \alpha_1)^{\beta_1} (x - \alpha_2)^{\beta_2} \cdots (x - \alpha_n)^{\beta_n}.$$

then

$$\begin{aligned} f'(x) &= \beta_1(x - \alpha_1)^{\beta_1 - 1} [(x - \alpha_2)^{\beta_2} \cdots (x - \alpha_n)^{\beta_n}] \\ &\quad + \beta_2(x - \alpha_2)^{\beta_2 - 1} [(x - \alpha_1)^{\beta_1} \cdots (x - \alpha_n)^{\beta_n}] + \cdots, \end{aligned}$$

Hence, for $x \neq \alpha_1, \alpha_2, \dots$

$$\frac{f'(x)}{f(x)} = \frac{\beta_1}{x - \alpha_1} + \frac{\beta_2}{x - \alpha_2} + \cdots + \frac{\beta_n}{x - \alpha_n}.$$

Thus,

$$\frac{a'_{2m}(r)}{a_{2m}(r)} = -2 \left(\frac{1}{2m + r} + \frac{1}{2m - 2 + r} + \cdots + \frac{1}{2 + r} \right),$$

or with $r = 0$

$$a'_{2m}(0) = -2 \left(\frac{1}{2m} + \frac{1}{2(m - 1)} + \cdots + \frac{1}{2} \right) a_{2m}(0).$$

Bessel's Equation Order Zero

Define

$$H_m = \frac{1}{m} + \frac{1}{m-1} + \cdots + \frac{1}{2} + 1,$$

then using this with the *recurrence relation*

$$a'_{2m}(0) = -H_m \frac{(-1)^m a_0}{2^{2m} (m!)^2}, \quad m = 1, 2, 3, \dots$$

It follows that the second solution of *Bessel's equation order zero* (with $a_0 = 1$) satisfies:

$$y_2(x) = J_0(x) \ln(x) + \sum_{m=1}^{\infty} \frac{(-1)^{m+1} H_m}{2^{2m} (m!)^2} x^{2m}, \quad x > 0.$$

Usually the second solution is taken to be the *Bessel function of the second kind of order zero*, which is defined as

$$Y_0(x) = \frac{2}{\pi} [y_2(x) + (\gamma - \ln(2))J_0(x)],$$


where γ is the Euler-Máscheroni constant

$$\gamma = \lim_{n \rightarrow \infty} (H_n - \ln(n)) \approx 0.5772.$$

Bessel's Equation Order Zero

9

The standard solutions for *Bessel's equation* are the *Bessel function of the first kind of order zero*, $J_0(x)$ and *Bessel function of the second kind of order zero*, $Y_0(x)$.



Roots Differing by an Integer, $r_1 - r_2 = N$

1

Case 3. Roots Differing by an Integer, $r_1 - r_2 = N$, where N is a positive integer.

As before, one solution of the *regular singular problem* satisfies:

$$y_1(x) = |x|^{r_1} \sum_{n=0}^{\infty} a_n x^n.$$

The second linearly independent solution has the form:

$$y_2(x) = k y_1(x) \ln |x| + |x|^{r_2} \sum_{n=0}^{\infty} b_n x^n$$

with these solutions converging for at least $|x| < \rho$, where ρ is the radius of convergence for $p(x)$ and $q(x)$.

This case divides into two subcases, depending on whether or not the logarithmic term appears, as k may be **zero**.

Roots Differing by an Integer, $r_1 - r_2 = N$

Case 3. Roots Differing by an Integer, $r_1 - r_2 = N$, where N is a positive integer.

This case is more complicated with the coefficients in the second solution satisfying:

$$b_n(r_2) = \frac{d}{dr} [(r - r_2)a_n(r)] \Big|_{r=r_2}, \quad n = 0, 1, 2, \dots$$

with $a_0 = r - r_2$ and

$$k = \lim_{r \rightarrow r_2} (r - r_2)a_N(r).$$

In practice the best way to determine if $k = 0$ is to compute $a_n(r_2)$ and see if one finds $a_N(r_2)$.

If this is possible, then the second solution is readily found without the logarithmic term; otherwise, the logarithmic term must be included.

Example: Roots, $r_1 - r_2 = N$

1

Example: Consider the *ODE*:

$$x^2 y'' + 3xy' + 4x^4 y = 0,$$

where $x = 0$ is a *regular singular point*.

We try a solution of the form: $y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r}$.

Differentiating y and entering into the equation gives:

$$\sum_{n=0}^{\infty} a_n (n+r)(n+r-1)x^{n+r} + 3 \sum_{n=0}^{\infty} a_n (n+r)x^{n+r} + 4 \sum_{n=4}^{\infty} a_{n-4} x^{n+r} = 0,$$

shifting the last index to match powers of x .

Examining this equation with $n = 0$ gives the *indicial equation*:

$$F(r) = r(r-1) + 3r = r(r+2) = 0,$$

which has the roots $r_1 = 0$ and $r_2 = -2$ ($r_1 - r_2 = 2$).

Example: Roots, $r_1 - r_2 = N$

2

Example: The series above could be rearranged in the following form:

$$\sum_{n=0}^{\infty} a_n(n+r)(n+r+2)x^{n+r} = -4 \sum_{n=4}^{\infty} a_{n-4}x^{n+r}.$$

With $r_1 = 0$, we obtain the *recurrence relations*:

$$a_1(r_1+1)(r_1+3) = 3a_1 = 0, \quad a_2(r_1+2)(r_1+4) = 8a_2 = 0, \quad a_3(r_1+3)(r_1+5) = 15a_3 = 0,$$

$a_1 = a_2 = a_3 = 0$, and

$$a_n(r) = -\frac{4}{(n+r)(n+r+2)}a_{n-4}(r) \quad \text{or} \quad a_n = -\frac{4}{n(n+2)}a_{n-4}.$$

It follows that

$$a_4 = -\frac{4}{6 \cdot 4}a_0 = -\frac{a_0}{3 \cdot 2}, \quad a_8 = -\frac{4}{10 \cdot 8}a_4 = \frac{a_0}{5!}, \quad a_{12} = -\frac{4}{14 \cdot 12}a_8 = -\frac{a_0}{7!},$$

and $a_1 = a_5 = \dots = a_{4n+1} = 0$, $a_2 = a_6 = \dots = a_{4n+2} = 0$, and
 $a_3 = a_7 = \dots = a_{4n+3} = 0$.

Example: Roots, $r_1 - r_2 = N$

Example: The results above are combined to give the **1st solution**:

$$y_1(x) = a_0 \sum_{m=0}^{\infty} \frac{(-1)^m}{(2m+1)!} x^{4m}.$$

To find the **2nd solution** we need to know if the **logarithmic term** needs to be included.

This term is unnecessary if

$$\lim_{r \rightarrow r_2} a_N(r) \text{ exists.}$$

For this example, $r_2 = -2$ and $N = 2$, so we examine

$$\lim_{r \rightarrow -2} a_2(r) = \frac{0}{(r+2)(r+4)} = 0,$$

which implies the second series may be computed directly with no **logarithmic term**.

Thus, we try a solution of the form:

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n+r_2},$$

Example: Roots, $r_1 - r_2 = N$

Example: Following the same process as finding $y_1(x)$, we have:

$$\sum_{n=0}^{\infty} b_n(n+r_2)(n+r_2+2)x^{n+r_2} = -4 \sum_{n=4}^{\infty} b_{n-4}x^{n+r_2}.$$

With $r_2 = -2$, we obtain the *recurrence relations*:

$$b_1(r_2+1)(r_2+3) = -b_1 = 0, \quad b_2(r_2+2)(r_2+4) = 0b_2 = 0, \quad b_3(r_2+3)(r_2+5) = 3b_3 = 0,$$

$b_1 = b_3 = 0$ (b_2 is arbitrary and generates y_1 , so take $b_2 = 0$), and

$$b_n(r) = -\frac{4}{(n+r)(n+r+2)}b_{n-4}(r) \quad \text{or} \quad b_n = -\frac{4}{n(n-2)}b_{n-4}.$$

It follows that

$$b_4 = -\frac{4}{4 \cdot 2}b_0 = -\frac{b_0}{2 \cdot 1}, \quad b_8 = -\frac{4}{8 \cdot 6}b_4 = \frac{b_0}{4!}, \quad b_{12} = -\frac{4}{12 \cdot 10}b_8 = -\frac{b_0}{6!},$$

and $b_1 = b_5 = \dots = b_{4n+1} = 0$, $b_2 = b_6 = \dots = b_{4n+2} = 0$, and
 $b_3 = b_7 = \dots = b_{4n+3} = 0$.

Example: Roots, $r_1 - r_2 = N$

Example: These results are combined to give the 2^{nd} solution:

$$y_2(x) = b_0 x^{-2} \sum_{m=0}^{\infty} \frac{(-1)^m}{(2m)!} x^{4m}.$$

It follows that our general solution for this example is:

$$y(x) = a_0 \sum_{m=0}^{\infty} \frac{(-1)^m}{(2m+1)!} x^{4m} + b_0 x^{-2} \sum_{m=0}^{\infty} \frac{(-1)^m}{(2m)!} x^{4m}.$$

This could be rewritten:

$$\begin{aligned} y(x) &= a_0 x^{-2} \sum_{m=0}^{\infty} \frac{(-1)^m}{(2m+1)!} (x^2)^{2m+1} + b_0 x^{-2} \sum_{m=0}^{\infty} \frac{(-1)^m}{(2m)!} (x^2)^{2m}, \\ &= x^{-2} (a_0 \sin(x^2) + b_0 \cos(x^2)). \end{aligned}$$

Example 2: Roots, $r_1 - r_2 = N$

1

Example 2: Consider the *ODE*:

$$x^2 y'' - xy = 0,$$

where $x = 0$ is a *regular singular point*.

Try a solution of the form: $y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r}$.

Differentiating y and entering into the equation gives:

$$\sum_{n=0}^{\infty} a_n (n+r)(n+r-1)x^{n+r} - \sum_{n=1}^{\infty} a_{n-1} x^{n+r} = 0,$$

shifting the last index to match powers of x .

Examining this equation with $n = 0$ gives the *indicial equation*:

$$F(r) = r(r-1) = 0,$$

which has the roots $r_1 = 1$ and $r_2 = 0$ ($r_1 - r_2 = 1$).

Example 2: Roots, $r_1 - r_2 = N$

2

Example 2: The series above is rearranged in the following form:

$$\sum_{n=0}^{\infty} a_n(n+r)(n+r-1)x^{n+r} = \sum_{n=1}^{\infty} a_{n-1}x^{n+r}.$$

With $r_1 = 1$, we obtain the *recurrence relation*:

$$a_n(r) = \frac{1}{(n+r)(n+r-1)}a_{n-1}(r) \quad \text{or} \quad a_n = \frac{1}{n(n+1)}a_{n-1}, \quad n = 1, 2, 3, \dots$$

It follows that

$$a_1 = \frac{1}{1 \cdot 2}a_0, \quad a_2 = \frac{1}{2 \cdot 3}a_1 = \frac{a_0}{2!3!}, \quad a_3 = \frac{1}{3 \cdot 4}a_2 = \frac{a_0}{3!4!},$$

so

$$a_n = \frac{a_0}{n!(n+1)!}, \quad n = 1, 2, 3, \dots$$

Example 2: Roots, $r_1 - r_2 = N$

Example 2: The results above are combined to give the **1st solution**:

$$y_1(x) = a_0 \sum_{n=0}^{\infty} \frac{1}{n!(n+1)!} x^{n+1}.$$

To find the **2nd solution** we need to know if the **logarithmic term** needs to be included.

This term is necessary if

$$\lim_{r \rightarrow r_2} a_N(r) \text{ fails to exist.}$$

For this example, $r_2 = 0$ and $N = 1$, so we examine

$$\lim_{r \rightarrow 0} a_1(r) = \frac{a_0(r)}{(r+1)r}.$$

Since a_0 is arbitrary (non-zero), this limit is undefined, so a second series solution requires the **logarithmic term**.

For $r_2 = 0$, we try a solution of the form:

$$y_2(x) = k y_1(x) \ln(x) + \sum_{n=0}^{\infty} b_n x^{n+r_2},$$

Example 2: Roots, $r_1 - r_2 = N$

Example 2: Insert $y_2(x) = k y_1(x) \ln(x) + \sum_{n=0}^{\infty} b_n x^n$ into the **ODE**, so

$$x^2 \left[k y_1'' \ln(x) + 2k y_1' \frac{1}{x} - k y_1 \frac{1}{x^2} + \sum_{n=2}^{\infty} n(n-1) b_n x^{n-2} \right] - k x y_1 \ln(x) - \sum_{n=0}^{\infty} b_n x^{n+1} = 0.$$

Because $y_1(x)$ is a solution of the **ODE**, $k \ln(x)[x^2 y_1'' - x y_1'] = 0$, which reduces this expression to

$$2k x y_1' - k y_1 + \sum_{n=2}^{\infty} n(n-1) b_n x^n - \sum_{n=0}^{\infty} b_n x^{n+1} = 0.$$

Using the series solution for $y_1(x)$ with $a_0 = 1$ and shifting indices, we obtain

$$\sum_{n=0}^{\infty} \left(\frac{2k}{(n!)^2} - \frac{k}{n!(n+1)!} \right) x^{n+1} + \sum_{n=1}^{\infty} b_{n+1} (n+1) n x^{n+1} - \sum_{n=0}^{\infty} b_n x^{n+1} = 0.$$

Example 2: Roots, $r_1 - r_2 = N$

Example 2: From the series above our *recurrence relation* gives:

$$k - b_0 = 0, \quad \text{or} \quad k = b_0$$

and

$$\frac{2k}{(n!)^2} - \frac{k}{n!(n+1)!} + n(n+1)b_{n+1} - b_n = 0, \quad n = 1, 2, 3, \dots$$

Equivalently,

$$b_{n+1} = \frac{1}{n(n+1)} \left[b_n - \frac{(2n+1)k}{n!(n+1)!} \right], \quad n = 1, 2, 3, \dots$$

For convenience we take $a_0 = 1$ and $b_0 = k = 1$.

The constant b_1 is still arbitrary (as it would generate $y_1(x)$ again), so we select $b_1 = 0$ and find a particular *2nd solution* using the *recurrence relation*:

$$y_2(x) = y_1(x) \ln(x) + 1 - \frac{3}{4}x^2 - \frac{7}{36}x^3 - \frac{35}{1728}x^4 - \dots$$