

Calculus for the Life Sciences

Lecture Notes – Integration by Substitution

Joseph M. Mahaffy,
(jmahaffy@mail.sdsu.edu)

Department of Mathematics and Statistics
Dynamical Systems Group
Computational Sciences Research Center
San Diego State University
San Diego, CA 92182-7720

<http://www-rohan.sdsu.edu/~jmahaffy>

Fall 2016

Outline

- 1 Introduction
- 2 Logistic Growth Model for Yeast
- 3 Integration by Substitution
 - Examples
- 4 Return to Logistic Growth
- 5 Examples
 - Integration by Substitution
 - Differential Equations
 - Logistic Growth
 - Lake Pollution with Seasonal Flow

Introduction

Introduction: Managing More Integrals

- To date we have learned a collection of basic integrals
 - Polynomials
 - Power Law
 - Exponentials - e^{kt}
 - Trig Functions - $\sin(kt)$ and $\cos(kt)$
- **Integration by substitution** allows a substitution that reduces the integral to a simpler form
- This is basically this inverse of the **Chain Rule of differentiation**
- Apply to models using separable differential equations
 - The logistic growth model
 - Model for motion of an object subject to gravity

Logistic Growth Model for Yeast

Logistic Growth Model for Yeast: Model considers a limited food source

- After a lag period, the organisms begin growing according to **Malthusian growth**
- As the food source becomes limiting, the growth of the organism slows and the population levels off
- This behavior is modeled by adding a negative quadratic term to the Malthusian growth model

$$\frac{dP}{dt} = rP \left(1 - \frac{P}{M} \right) \quad \text{with} \quad P(0) = P_0$$

Logistic Growth Model for Yeast

Experiment: G. F. Gause (*Struggle for Existence*) studied standard brewers yeast, *Saccharomyces cerevisiae*


- *S. cerevisiae* placed in a closed vessel, where nutrient was changed regularly (every 3 hours)
- Simulates a constant source of nutrient

Time (hr)	0	1.5	9	10	18	18	23
Volume	0.37	1.63	6.2	8.87	10.66	10.97	12.5
Time (hr)	25.5	27	34	38	42	45.5	47
Volume	12.6	12.9	13.27	12.77	12.87	12.9	12.7

Logistic Growth Model for Yeast

3

Graph of data and best fitting model



Logistic Growth Model for Yeast

4

Model: The **Logistic Growth Model** that best fits the data is

$$\frac{dP}{dt} = 0.259 P \left(1 - \frac{P}{12.7} \right), \quad \text{with } P(0) = 1.23$$

- How do we find the solution to this nonlinear differential equation?
- This is a separable equation
- The integral for P involves two integration techniques
- We'll concentrate on the the **integration by substitution**

Integration by Substitution

Integration by Substitution

- Integration is the inverse of differentiation
- Many functions that do not have an antiderivative
- **Integration by substitution** extends the number of integrable functions
- This technique is the inverse of the chain rule of differentiation
- The substitution technique finds a function that reduces an integral to an easier form

Example 1

Example 1: Let a be a constant and consider the integral

$$\int (x + a)^n dx$$

Make the substitution $u = x + a$, and the derivative gives the differentials $du = dx$, so

$$\begin{aligned}\int (x + a)^n dx &= \int u^n du \\ &= \frac{u^{n+1}}{n+1} + C \\ &= \frac{(x + a)^{n+1}}{n+1} + C\end{aligned}$$

Example 2

Example 2: Consider the integral

$$\int x e^{-x^2} dx$$

Make the substitution $u = -x^2$, and the derivative gives the differentials $du = -2x dx$, so

$$\begin{aligned}\int x e^{-x^2} dx &= \int e^{-x^2} \left(-\frac{1}{2}\right) (-2x) dx \\ &= -\frac{1}{2} \int e^u du \\ &= -\frac{1}{2} e^u + C \\ &= -\frac{1}{2} e^{-x^2} + C\end{aligned}$$

Example 3

Example 3: Consider the integral

$$\int (x^2 + 2x + 4)^3 (x + 1) dx$$

Make the substitution $u = x^2 + 2x + 4$, and the derivative gives the differentials $du = (2x + 2)dx$, so

$$\begin{aligned} \int (x^2 + 2x + 4)^3 (x + 1) dx &= \frac{1}{2} \int (x^2 + 2x + 4)^3 (2x + 2) dx \\ &= \frac{1}{2} \int u^3 du \\ &= \frac{u^4}{8} + C \\ &= \frac{(x^2 + 2x + 4)^4}{8} + C \end{aligned}$$

Integration by Substitution

Integration by Substitution: What makes a good substitution?

- Choose u such that when u and du are substituted for the expression of x under the integrand, the remaining integral became one of the basic integrals solved earlier
- There are a few choices that are very natural for a substitution
 - Let u be any expression of x in the exponent of the exponential function e or the argument of any trigonometric functions or the logarithm function
 - Let u be an expression of x inside parentheses raised to a power, where you should be able to see the derivative of that expression multiplying this expression to a power

Return to Logistic Growth

Return to Logistic Growth: The **Logistic Growth Model** is

$$\frac{dP}{dt} = rP \left(1 - \frac{P}{M}\right) = -rP \left(\frac{P}{M} - 1\right)$$

- Separate Variables to give

$$\int \frac{dP}{P \left(\frac{P}{M} - 1\right)} = -r \int dt$$

- The integral on the right is very easy to solve
- The integral on the left requires a technique from algebra
 - Fraction is split into two simple fractions (reverse of a common denominator)

$$\frac{1}{P \left(\frac{P}{M} - 1\right)} = \frac{\frac{1}{M}}{\left(\frac{P}{M} - 1\right)} - \frac{1}{P}$$

Return to Logistic Growth

2

Separated Differential Equation: From fractional form above, write the integral as

$$\int \frac{dP}{P \left(\frac{P}{M} - 1\right)} = \frac{1}{M} \int \frac{dP}{\left(\frac{P}{M} - 1\right)} - \int \frac{dP}{P}$$

- One integral is easy

$$\int \frac{dP}{P} = \ln |P| + C$$

- For the other make the substitution $u = \frac{P}{M} - 1$, so $du = \frac{dP}{M}$

$$\frac{1}{M} \int \frac{dP}{\left(\frac{P}{M} - 1\right)} = \int \frac{du}{u} = \ln |u| = \ln \left| \frac{P}{M} - 1 \right|$$

Return to Logistic Growth

Separated Differential Equation:

$$\int \frac{dP}{P\left(\frac{P}{M} - 1\right)} = -r \int dt = -rt + C$$

- From results above

$$\ln \left| \frac{P}{M} - 1 \right| - \ln |P| = -rt + C$$

- Thus,

$$\ln \left| \frac{\frac{P}{M} - 1}{P} \right| = \ln \left| \frac{P - M}{MP} \right| = -rt + C$$

- Exponentiating,

$$\left| \frac{P(t) - M}{MP(t)} \right| = e^{-rt+C}$$

Return to Logistic Growth

Solution: Removing the absolute value

$$\frac{P(t) - M}{MP(t)} = Ae^{-rt}$$

- Solving for $P(t)$ gives

$$P(t) = \frac{M}{1 - MAe^{-rt}}$$

- With the initial condition, $P(0) = P_0$

$$P_0 = \frac{M}{1 - MA} \quad \text{or} \quad A = \frac{P_0 - M}{MP_0}$$

- Inserting this into the solution above gives

$$P(t) = \frac{P_0M}{P_0 + (M - P_0)e^{-rt}}$$

Return to Logistic Growth

Yeast Model: The best fitting yeast model

$$\frac{dP}{dt} = 0.259 P \left(1 - \frac{P}{12.7} \right), \quad \text{with } P(0) = 1.23$$

- The general logistic solution is

$$P(t) = \frac{P_0 M}{P_0 + (M - P_0) e^{-rt}}$$

- It follows that

$$P(t) = \frac{15.62}{1.23 + 11.47 e^{-0.259t}}$$

- This function creates the standard **S-shaped curve** of logistic growth and has the **carrying capacity** of **12.7**

Integration Example 1

Integration Example 1: Consider the integral

$$\int x^2 \cos(4 - x^3) dx$$

Skip Example

Solution: A natural substitution is

$$u = 4 - x^3 \quad \text{so} \quad du = -3x^2 dx$$

The solution of the integral is

$$\begin{aligned} \int x^2 \cos(4 - x^3) dx &= -\frac{1}{3} \int \cos(4 - x^3) (-3x^2) dx \\ &= -\frac{1}{3} \int \cos(u) du \\ &= -\frac{1}{3} \sin(u) + C \\ &= -\frac{1}{3} \sin(4 - x^3) + C \end{aligned}$$

Integration Example 2

Integration Example 2: Consider the integral

$$\int \frac{(\ln(2x))^2}{x} dx$$

Skip Example

Solution: A natural substitution is

$$u = \ln(2x) \quad \text{so} \quad du = \frac{dx}{x}$$

The solution of the integral is

$$\begin{aligned} \int \frac{(\ln(2x))^2}{x} dx &= \int u^2 du \\ &= \frac{u^3}{3} + C \\ &= \frac{1}{3} (\ln(2x))^3 + C \end{aligned}$$

Differential Equation Example 1

1

Differential Equation Example 1: Consider

$$\frac{dy}{dt} = \frac{2ty}{t^2 + 4}, \quad y(0) = 8$$

Skip Example

Solution: Separate the differential equation into the two integrals

$$\int \frac{dy}{y} = \int \frac{2t}{t^2 + 4} dt$$

The right integral uses the substitution $u = t^2 + 4$, so $du = 2t dt$

$$\ln |y(t)| = \int \frac{du}{u} = \ln |u| + C = \ln(t^2 + 4) + C$$

Differential Equation Example 1

2

Solution (cont): The integrations give

$$\ln |y(t)| = \ln(t^2 + 4) + C$$

- Exponentiating

$$y(t) = e^{\ln(t^2+4)+C} = e^C(t^2 + 4)$$

- Note that e^C could be positive or negative depending on the initial condition
- From the initial condition, $y(0) = 8$, it follows that

$$y(t) = 2(t^2 + 4)$$

Differential Equation Example 2

1

Differential Equation Example 2: Consider

$$\frac{dy}{dt} = 2t e^{t^2-y}, \quad y(0) = 2$$

Skip Example

Solution: Rewrite the differential equation

$$\frac{dy}{dt} = 2t e^{t^2} e^{-y}$$

Separate the differential equation into the two integrals

$$\int e^y dy = \int 2t e^{t^2} dt$$

Differential Equation Example 2

2

Solution (cont): The right integral uses the substitution $u = t^2$, so $du = 2t dt$

$$\int e^y dy = e^y = \int 2t e^{t^2} dt = \int e^u du = e^u + C$$

- By substitution the implicit solution is

$$e^y = e^{t^2} + C$$

- Taking logarithms

$$y(t) = \ln(e^{t^2} + C)$$

- From the initial condition, $y(0) = 2 = \ln(1 + C)$, it follows that

$$y(t) = \ln(e^{t^2} + e^2 - 1)$$

Logistic Growth

1

Logistic Growth: Suppose that a population of animals satisfies the logistic growth equation

$$\frac{dP}{dt} = 0.01 P \left(1 - \frac{P}{2000} \right), \quad P(0) = 50$$

- Find the general solution of this equation
- Determine how long it takes for this population to double
- Find how long it takes to reach half of the carrying capacity

Logistic Growth

2

Solution: We separate this logistic growth model

$$\int \frac{dP}{P \left(\frac{P}{2000} - 1 \right)} = -0.01 \int dt = -0.01 t + C$$

- The **Fundamental Theorem Algebra** gives

$$\frac{1}{P \left(\frac{P}{2000} - 1 \right)} = \frac{\frac{1}{2000}}{\left(\frac{P}{2000} - 1 \right)} - \frac{1}{P}$$

- We use the substitution $u = \frac{P}{2000} - 1$, so $du = \frac{dP}{2000}$

$$\frac{1}{2000} \int \frac{dP}{\left(\frac{P}{2000} - 1 \right)} - \int \frac{dP}{P} = \int \frac{du}{u} - \int \frac{dP}{P} = -0.01 t + C$$

Logistic Growth

3

Solution (cont): From the substitution $u = \frac{P}{2000}$

$$\int \frac{du}{u} - \int \frac{dP}{P} = -0.01t + C$$

- Thus,

$$\ln |u| - \ln |P| = \ln \left| \frac{P - 2000}{2000} \right| - \ln |P| = -0.01t + C$$

- So,

$$\ln \left| \frac{P - 2000}{2000 P} \right| = -0.01t + C$$

Logistic Growth

4

Solution (cont): Exponentiating the previous expression

$$\frac{P(t) - 2000}{2000 P(t)} = e^{-0.01 t + C} = A e^{-0.01 t}$$

- Solving for $P(t)$,

$$P(t) = \frac{2000}{1 - 2000A e^{-0.01 t}}$$

- With the initial condition, $P(0) = 50$,

$$P(t) = \frac{2000}{1 + 39 e^{-0.01 t}}$$

Logistic Growth

Solution (cont): The logistic growth model is

$$P(t) = \frac{2000}{1 + 39e^{-0.01t}}$$

- The population doubles when

$$P(t_d) = \frac{2000}{1 + 39e^{-0.01t_d}} = 100$$

- Thus,

$$1 + 39e^{-0.01t_d} = 20 \quad \text{or} \quad e^{0.01t_d} = \frac{39}{19}$$

- Solving for doubling time

$$t_d = 100 \ln\left(\frac{39}{19}\right) = 71.9$$

Logistic Growth

Solution (cont): The logistic growth model is

$$P(t) = \frac{2000}{1 + 39e^{-0.01t}}$$

- The population reaches half the carrying capacity when

$$P(t_h) = \frac{2000}{1 + 39e^{-0.01t_h}} = 1000$$

- Thus,

$$1 + 39e^{-0.01t_h} = 2 \quad \text{or} \quad e^{0.01t_h} = 39$$

- Solving for doubling time

$$t_h = 100 \ln(39) = 366.4$$

Lake Pollution with Seasonal Flow

1

Lake Pollution with Seasonal Flow Often the flow rate into a lake varies with the season

- Suppose that a 200,000 m³ lake maintains a constant volume and is initially clean
- A river flowing into the lake has 6 μg/m³ of a pesticide
- Assume that the flow of the river has the sinusoidal form

$$f(t) = 100(2 - \cos(0.0172t)),$$

where t is in days

- Find and solve the differential equation describing the concentration of the pesticide in the lake
- Graph the solution for 2 years

Lake Pollution with Seasonal Flow

2

Solution: Begin by creating the differential equation

- The change in the amount of pesticide, $A(t)$, equals the amount entering - the amount leaving

$$\frac{dA(t)}{dt} = 600(2 - \cos(0.0172t)) - 100(2 - \cos(0.0172t))c(t)$$

- Concentration satisfies $c(t) = \frac{A(t)}{200,000}$, so

$$\frac{dc}{dt} = -\frac{(2 - \cos(0.0172t))}{2000}(c - 6)$$

- Separating variables

$$\int \frac{dc}{c - 6} = -\frac{1}{2000} \int (2 - \cos(0.0172t))dt$$

Lake Pollution with Seasonal Flow

Solution: By letting $u = c - 6$ with $du = dc$, the integrals are

$$\int \frac{du}{u} = -0.0005 \int (2 - \cos(0.0172t)) dt$$

- Integrating

$$\ln(u) = \ln(c(t) - 6) = -0.0005 \left(2t - \frac{\sin(0.0172t)}{0.0172} \right) + C$$

- By exponentiating this implicit solution, using the initial condition ($c(0) = 0$), and letting $\frac{1}{0.0172} = 58.14$, the solution becomes

$$c(t) = 6 \left(1 - e^{-0.0005(2t - 58.14 \sin(0.0172t))} \right)$$

Lake Pollution with Seasonal Flow

4

Graph: Consider solution for 2 yr or 730 days

$$c(t) = 6 \left(1 - e^{-0.0005(2t - 58.14 \sin(0.0172t))} \right)$$

