

HW 6

5.5.1. A Sturm-Liouville eigenvalue problem is called self-adjoint if

$$p \left(u \frac{dv}{dx} - v \frac{du}{dx} \right) \Big|_a^b = 0$$

10

since then $\int_a^b [uL(v) - vL(u)] dx = 0$ for any two functions u and v satisfying the boundary conditions. Show that the following yield self-adjoint problems.

- (c) $\frac{d\phi}{dx}(0) - h\phi(0) = 0$ and $\frac{d\phi}{dx}(L) = 0$
- (d) $\phi(a) = \phi(b)$ and $p(a)\frac{d\phi}{dx}(a) = p(b)\frac{d\phi}{dx}(b)$

5.5.5. Consider

$$L = \frac{d^2}{dx^2} + 6\frac{d}{dx} + 9.$$

10

- (a) Show that $L(e^{rx}) = (r+3)^2 e^{rx}$.
- (b) Use part (a) to obtain solutions of $L(y) = 0$ (a second-order constant-coefficient differential equation).
- (c) If z depends on x and a parameter r , show that

$$\frac{\partial}{\partial r} L(z) = L \left(\frac{\partial z}{\partial r} \right).$$

- (d) Using part (c), evaluate $L(\partial z / \partial r)$ if $z = e^{rx}$.
- (e) Obtain a second solution of $L(y) = 0$, using part (d).

5.5.8. Consider a fourth-order linear differential operator,

15

$$L = \frac{d^4}{dx^4}.$$

- (a) Show that $uL(v) - vL(u)$ is an exact differential.
- (b) Evaluate $\int_0^1 [uL(v) - vL(u)] dx$ in terms of the boundary data for any functions u and v .
- (c) Show that $\int_0^1 [uL(v) - vL(u)] dx = 0$ if u and v are any two functions satisfying the boundary conditions

$$\begin{aligned} \phi(0) &= 0 & \phi(1) &= 0 \\ \frac{d\phi}{dx}(0) &= 0 & \frac{d^2\phi}{dx^2}(1) &= 0. \end{aligned}$$

- (d) Give another example of boundary conditions such that

$$\int_0^1 [uL(v) - vL(u)] dx = 0.$$

- (e) For the eigenvalue problem [using the boundary conditions in part (c)]

$$\frac{d^4\phi}{dx^4} + \lambda e^x \phi = 0,$$

show that the eigenfunctions corresponding to different eigenvalues are orthogonal. What is the weighting function?

HW 6 (cont.)

5.5.11. *(a) Suppose that

15

$$L = p(x) \frac{d^2}{dx^2} + r(x) \frac{d}{dx} + q(x).$$

Consider

$$\int_a^b vL(u) dx.$$

By repeated integration by parts, determine the adjoint operator L^* such that

$$\int_a^b [uL^*(v) - vL(u)] dx = H(x) \Big|_a^b.$$

What is $H(x)$? Under what conditions does $L = L^*$, the self-adjoint case? [Hint: Show that

$$L^* = p \frac{d^2}{dx^2} + \left(2 \frac{dp}{dx} - r \right) \frac{d}{dx} + \left(\frac{d^2 p}{dx^2} - \frac{dr}{dx} + q \right)].$$

(b) If

$$u(0) = 0 \quad \text{and} \quad \frac{du}{dx}(L) + u(L) = 0,$$

what boundary conditions should $v(x)$ satisfy for $H(x) \Big|_0^L = 0$, called the adjoint boundary conditions?

20

5.8.5. Consider

$$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2}$$

with $\frac{\partial u}{\partial x}(0, t) = 0$, $\frac{\partial u}{\partial x}(L, t) = -hu(L, t)$, and $u(x, 0) = f(x)$.

- 8 (a) Solve if $h > 0$.
- 12 (b) Solve if $h < 0$.

25

5.8.8. Consider the boundary value problem

$$\frac{d^2 \phi}{dx^2} + \lambda \phi = 0 \quad \text{with} \quad \begin{aligned} \phi(0) - \frac{d\phi}{dx}(0) &= 0 \\ \phi(1) + \frac{d\phi}{dx}(1) &= 0. \end{aligned}$$

- 5 (a) Using the Rayleigh quotient, show that $\lambda \geq 0$. Why is $\lambda > 0$?
- 5 (b) Prove that eigenfunctions corresponding to different eigenvalues are orthogonal.
- 10 *(c) Show that

$$\tan \sqrt{\lambda} = \frac{2\sqrt{\lambda}}{\lambda - 1}.$$

Determine the eigenvalues graphically. Estimate the large eigenvalues.

HW 6 (cont.)

5 (d) Solve

$$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2}$$

with

$$u(0, t) - \frac{\partial u}{\partial x}(0, t) = 0$$

$$u(1, t) + \frac{\partial u}{\partial x}(1, t) = 0$$

$$u(x, 0) = f(x).$$

You may call the relevant eigenfunctions $\phi_n(x)$ and assume that they are known.

5

5.8.11. Determine all negative eigenvalues for

$$\frac{d^2 \phi}{dx^2} + 5\phi = -\lambda\phi \text{ with } \phi(0) = 0 \text{ and } \phi(\pi) = 0.$$